Stability in Matching with Externalities: Pairs Competition and Oligopolistic Joint Ventures

Kenzo Imamura

1 Introduction

matchings might be realized by forming a pair, and showed the existence of stable matching under pessimism as in Sasaki and Toda (1996). Chen (2019) considered a speci...c example of Cournot oligopoly game played by joint ventures, and assumed that every potential partner for a company induces a unique consistent expectation for the realized matching. With this list of expectations for each possible pair, stable matching is de...ned as the outcome of this game. Chen identi..ed conditions under which positively and negatively assortative matchings are stable.

In these papers, each player has expectations on the realization of a matching when she is partnered with each of the players on the other side of the market, and stable matching is built on these expectations. In contrast, this paper mirrors the original de...nition of pairwise stable matching in matching problems without externalities. Our pairwise stability starts with a matching and checks whether or not there is a pair of players with a pro...table deviation away from the original matching. There is, however, a subtle issue in the presence of externalities— the rest of matching matters. Thus, we need to formulate the matching induced by the deviation of a pair from the original matching. We specify this using an exectiveness function, and consider two exectiveness functions speci...cally. The ..rst is that after a deviation by a pair, the dumped partners stay single and no other player changes their partners. The second is that after a deviation by a pair, the dumped partners form a pair and all other players remain with their partners. The former exectiveness function is in the literature of theory of coalition formation, and is adopted to analyze convergence of a sequence of myopic deviations in a marriage problem by Roth and Vande Vate (1990). The latter exectiveness function was proposed by Knuth (1976) in the context of the marriage problem in which all players are acceptable to all other players.

To see the di¤erence between these two e¤ectiveness functions, consider the example of a pairs ...gure skating competition. Suppose that there are three male and three female skaters, one with high, medium, and low ability in each gender. Moreover, suppose that there are complementarities in partners' abilities. Then, it is natural for them to have a (positively) assortative matching, since a high ability partner is always desirable. However, this assortative matching may not be pairwise stable in the Roth-Vande Vate sense. Consider a deviation by the high ability male and the

1

swapping. In Section 4, we consider an oligopolistic joint ventures problem, which is an assignment game version of the pairs competition model without endogenous exorts. We show that pairwise stable assignments can be supported only by the assortative matching, and characterize the one-side optimal stable assignment. In Section 5, we introduce personalized intrinsic utility from partners and heterogeneous match qualities of players, and show that pairwise stable assignment via swapping may not exist. Section 6 concludes.

1.1 A Brief Literature Review

There are three branches of literature that are related to the current paper. The ...rst branch is the one of matching with externalities. Recently, a number of papers have been written in this ...eld in addition to Sasaki and Toda (1996), Hafalir (2008), and Chen (2019). Mumcu and Saglam (2010), Fisher and Hafalir (2016), and Chade and Eeckout (2020) all dealt with one-to-one matching problems with externalities in di¤erent ways. Mumcu and Saglam (2010) introduced outside options, and Fisher and Hafalir (2016) and Chade and Eeckhout (2020) removed the impacts of pairwise deviations through externalities by imposing a behavioral assumption and by considering a continuum of atomless agents, respectively. Bando (2012, 2014), and Pycia and

```
( ^{0}( (m)) = (m)), and if w is paired under ( (w) \in w) then w's partner (w) is single under ^{0} ( ^{0}( (w)) = (w)); and (iii) for all
```

($(m_i)\ 2\ W$), team i's members' exorts are aggregated by a CES function

$$Y_i = (a_m e_m + a_{(m)} e_{(m)})^{\frac{1}{2}};$$
 (1)

where

and player x's equilibrium exort given Yi and Y can be written as

$$e_x = Y_i (1 i) \frac{1}{Y} a_x^{\frac{1}{1-}} a_x^{\frac{1}{1-}}$$
 (3)

Raising this to the power of and then multiply it by a_x ,

$$a_x e_x = Y_i$$
 $(1 \quad _i) \frac{1}{Y}$ $^{\frac{1}{1-}} a_x^{\frac{1}{1-}} V^{\frac{1}{1-}}$

is obtained (the power of a_x is calculated by $\frac{2}{1} + \frac{2}{1}$). Substituting this back to (1), we obtain

$$Y_i = Y_i (1 i) \frac{1}{Y} a_x^{\frac{1}{1-}} + a_x^{\frac{1}{1-}} V_x^{\frac{1}{1-}}$$

or

$$\frac{1}{A_i()} = \frac{Y_i}{Y^2} V; \tag{4}$$

where $A_i() = a_m^{\frac{1}{1-}} + a_{(m)}^{\frac{1}{1-}}$ stands for the

stable if and only if (i) R_m or R_w for any pairwise deviations (m; w) 2 M W with ! (m;w) , and (ii) R_x for any single player deviation x 2 M [W with ! $_x$. The following example shows that there may not be a pairwise stable matching.

Example 1. Consider a ...gure skating contest with $M = fm_1$; m_2 ; m_3g and $W = fw_1$; w_2 ; w_3g . Let $= \frac{1}{2}$, $a_{m_1} = a_{w_1} = 1$, $a_{m_2} = a_{w_2} = 0.9$, and $a_{m_3} = a_{w_3} = 0.7$. We calculate m_1 's payows under the assortative matching and the one after he deviates with w_2 :

(i) =
$$f(m_1; w_1); (m_2; w_2); (m_3; w_3)g$$
:

$$U_{m_1}() = 1 \frac{2 \frac{1}{2}}{\frac{1}{2} + \frac{1}{1 \cdot 8} + \frac{1}{1 \cdot 4}} 1 \frac{2 \frac{1}{2}}{\frac{1}{2} + \frac{1}{1 \cdot 8} + \frac{1}{1 \cdot 4}} \frac{1}{2} = 0:31209$$

(ii) ${}^{\emptyset} f(m_1; w_2); (m_3; w_3)g:$

$$U_{m_1}(^{0}) = 1 \quad \frac{\frac{1}{1:9}}{\frac{1}{1:9} + \frac{1}{1:4}} \quad 1 \quad \frac{\frac{1}{1:9}}{\frac{1}{1:9} + \frac{1}{1:4}} \quad \frac{1}{1:9} = 0:44720$$

Thus, agent m_1 is better o^x by dumping his higher ability partner for an inferior partner. A similar deviation blocks any other fully matched matching, and if agents are not fully matched in matching , then is blocked by an unmatched pair. Thus, there is no pairwise stable matching in this example.

The problem underlying this example is that players prefer to have a smaller number of rival pairs, and the best player would rather have a weaker partner if the number of rival pairs goes down. However, since single players cannot participate in the competition, resulting in receiving the lowest payo $^{\alpha}$ s, it does not make sense to expect that they will stay singles. If the single players becomes a pair, the number of rivals do not change, undermining the motivation for the best player to seek a lower ability partner. Using the second exectiveness function \Rightarrow_S allows us to de..ne the following alternative stability concept. A matching is pairwise stable \forall ia swapping if and only if (i) R_m or R_w for any pairwise deviations (m; w) 2 M W with $\Rightarrow_{(m:w)}$. In the following, we will show that the assortative matching is uniquely stable in the above sense. We ..rst prove the following lemma, which demonstrates that an assortative swapping improves higher ability players' payo $^{\alpha}$ s.

Lemma 1. Let ,

Now, we borrow the model from Shubik (1984) to describe our oligopolistic market.⁷ Suppose that there are n products produced by n active joint ventures together with a numeraire commodity (the 0th commodity). There is unit mass of identical consumers, each with a quadratic utility function:

$$u = \sum_{i=1}^{N} x_i \frac{1}{2} \sum_{i=1}^{N} x_i^2 \frac{X^2}{2} \sum_{j=1}^{N} x_k x_j + x_0;$$
 (5)

where 2 [0; 1) is a substitution parameter between products. As increases, substitutability

By Lemmas 3 and 4, we know that X is strictly supermodular and strictly increasing. The following proposition shows that there are pairwise stable assignments, and characterizes the M-optimal pairwise stable matching by using the above output matrix X.⁸

Proposition 5. In an oligopolistic joint venture model, there exist pairwise stable assignments. Under the M-optimal pairwise stable assignment, the pairwise stable payo¤ vector for W is minimized at $s=(s_1; ...; s_n)$ where $s_j=\frac{P_{\substack{n=1\\j'=j}}^n(X_{j'+1j'}-X_{j'+1j'+1})}{j'=j}$ for any j=1 and $s_n=0$, and the stable payo¤ vector for M is calculated by $r_j=X_{j,j}-s_j$.

Bulow and Levin (2006) derived the above simple "minimum competitive salary" formula in the context of ...rm-worker matching problem with output function $X_{ij}=a_i-a_j$ (thus with

$$U_{x}() = b_{x}((x)) + (1) U_{x}()$$

where $U_x(\)$ is the same as in Section 3:

$$U_{x}(\)=\ 1 \quad \frac{(n(\)\ 1)\frac{1}{A(\)}}{\sum_{j=1}^{n}\frac{1}{A(\)}}^{\#"} \quad 1 \quad \frac{(n(\)\ 1)\frac{1}{A(\)}}{\sum_{j=1}^{n}\frac{1}{A(\)}} \quad \frac{a_{x}}{A_{i}(\)} \quad ^{\frac{\#}{1-}}V:$$

Clearly, when = 0, this problem degenerates to the pairs competition problem, and to the standard one-to-one matching problem without externalities when = 1. In the neighborhoods of = 1 or = 0, pairwise stable matching via swapping obviously exists. But does this hold when is signi...cantly far from the end points? Unfortunately, in general, pairwise stable matching via swapping may not exist as we can see from the following example.

Example 2. Consider a ...gure skating contest with $M=fm_1;m_2;m_3g$ and $W=fw_1;w_2;w_3g$. Let $=\frac{1}{2}$, $a_{m_1}=a_{w_1}=0.5$, $a_{m_2}=a_{w_2}=0.4$, and $a_{m_3}=a_{w_3}=0.25$. Personal intrinsic payoxs from the partners are described by the following matrices:

- (ii) $^2 = f(m_1; w_1); (m_2; w_3); (m_3; w_2)g: U_{m_1}(^2) = U_{w_1}(^2) = 0.384; U_{m_2}(^2) = U_{w_2}(^2) = 0.131; U_{m_3}(^2) = U_{w_3}(^2) = 0.174.$
- (iii) $^3 = f(m_1; w_3); (m_2; w_1); (m_3; w_2)g; U_{m_1}(^3) = 0:183; U_{m_2}(^3) = 0:332; U_{m_3}(^3) = 0:160; U_{w_1}(^3) = 0:305; U_{w_2}(^3) = 0:119; U_{w_3}(^3) = 0:257.$

 m_2 , and m_2 to m_1 .¹⁰ Therefore, it is not easy to assure the existence of pairwise stable matching via swapping. If, however, ability ranking agrees with intrinsic preference ranking for all agents, then we have the following result.

Proposition 6. Suppose that b_m (w_1) b_m (w_2) ::: b_m (w_n) for all m_i 2 M and b_w (m_1) b_w (m_2) ::: b_w (m_n) for all w_i 2 W. Then, is the unique pairwise stable matching via swapping.

5.2 Match Qualities

Here, we introduce match qualities of pairs, and we introduce match qualities in the oligopolistic joint venture problem. Let a match quality matrix Q be

where q_{ij} 2 (0;1] which captures how good a match between m_i and w_j is. Exectively, this describes how well m_i and w_i can work together.

In this case, the structure of the problem are the same as the original oligopolistic joint ventures, so our solution concept, the pairwise stable assignment via swapping, is well de..ned. However, with an arbitrary match quality matrix Q, a pairwise stable assignment via swapping may not exist. This is because we can create an arbitrary matrix of the marginal cost of each joint venture, $C = (c(m_i; w_j))_{i:j=1;...;n}$ by freely choosing Q. The following example demonstrates this result.

Example 3. Let $M = fm_1$; m_2 ; m_3g and $W = fw_1$; w_2 ; w_3g with

$$C = \begin{cases} 0.1 & 0.11 & 0.3 \\ 0.3 & 0.1 & 0.3 \\ 0.3 & 0.3 & 0.3 \end{cases}$$

There are six full matchings:

(i)
$$_{1}(m_{1}) = w_{1}$$
; $_{1}(m_{2}) = w_{2}$; $_{1}(m_{3}) = w_{3}$; $(m_{1}; w_{2})$ deviates to create $_{2}$.

(ii)
$$_{2}(m_{1}) = w_{2}$$
; $_{2}(m_{2}) = w_{1}$; $_{1}(m_{3}) = w_{3}$; $(m_{2}; w_{3})$ deviates to create $_{3}$.

(iii)
$$_{3}(m_{1}) = w_{2}; _{3}(m_{2}) = w_{3}; _{3}(m_{3}) = w_{1}; (m_{1}; w_{1})$$
 deviates to create $_{4}$.

Proposition 7. In the oligopolistic competition by joint ventures model, there exists a pairwise stable matching via swapping if = 0 (no externalities: local monopoly).

6 Concluding Remarks

This implies that Y_i is

$$Y_i = {}_{i}Y = {}^{i}1$$
 $\frac{(n())}{\sum_{j=1}^{n} \frac{1}{A(j)}} {}^{\#} {}^{"} \underbrace{(n())}_{j=1} {}^{1}\underbrace{\frac{1}{A(j)}}_{A(j)} {}^{\#} V$

These results lead to the following formulas that are essential for the analysis of stability of team structure. Recalling (3), we obtain

This implies that agent i's payo¤ is written as

In order to show that Regularity Condition 1 assures $_{i}>0$ for all i=1;:::;n, we use Lemma 1 (i) below. Repeatedly applying Lemma 1 (i), it is easy to see $P_{j=1}^{n} \frac{1}{A(*)} P_{j=1}^{n} \frac{1}{A()}$ for all . Moreover, $A_{n}()$ $A_{i}()$ for all . Thus, if the Regularity condition is satis...ed, then $_{i}()>0$ for all and all i=1;:::;n. We have completed the proof.

Proof of Lemma 1. Let $A_j()$ $a(m_j)^{-} + a((m_j))^{-}$

Thus,

$$\frac{1}{A^{\cdot}() + \tilde{A}_{k}()} + \frac{1}{A_{k}()} + \frac{1}{A_{k}()} + \frac{1}{A_{k}()} + \frac{1}{A_{k}()} = A^{\cdot}() + \tilde{A}_{k}() + \tilde{A$$

Summing up over commodities produces:

$$n \qquad \begin{matrix} x_1 \\ x_j \end{matrix} \qquad \begin{matrix} x_1 \\ x_j \end{matrix} \qquad \begin{matrix} x_1 \\ x_j \end{matrix} = \begin{matrix} x_1 \\ x_j \end{matrix} = \begin{matrix} x_1 \\ x_j \end{matrix} ;$$

or

$$X^{n}$$
 $x_{j} = \frac{1}{1+n}(n - P);$

where $P = P_{j=1}^{n} p_{j}$. Substituting this back to the f.o.c., we obtain:

$$x_i = \frac{1+n}{1+n}(n - P) = p_i;$$

or

$$x_i = x_i(p_i; P) = p_i \frac{1}{1+p}(n P)$$
:

Thus, the market demand function for good i is:

$$x_i(p_i; P) = \frac{1}{1+n} + \frac{1}{1+n} P \quad p_i$$
:

We have completed the proof.

■

Proof of Proposition 3. The ..rm i's f.o.c. with respect to p_i is:

$$\frac{1}{1+n} + \frac{1}{1+n}P$$
 $p_i + (p_i c_i) \frac{1}{1+n}$ $1 = 0;$

or

$$1 + \frac{1 + (n - 1)}{1 + n}$$
 $p_i = \frac{1 + n}{1 + n} + \frac{1 + (n - 1)}{1 + n} c_i$

or

$$\frac{2 + (2n - 1)}{1 + n} p_i = \frac{1 + n}{1 + n} + \frac{1 + (n - 1)}{1 + n} c_i$$

$$p_i = \frac{1 + (n - 1)}{2 + (2n - 1)} + \frac{1 + (n - 1)}{2 + (2n - 1)} C_i$$

Summing them up, we have

$$P = \frac{n}{2 + (2n - 1)} + \frac{n}{2 + (2n - 1)}P + \frac{1 + (n - 1)}{2 + (2n - 1)}X^{n}$$

Thus,

$$\frac{2 + (n - 1)}{2 + (2n - 1)}P = \frac{n}{2 + (2n - 1)} + \frac{1 + (n - 1)}{2 + (2n - 1)} X^{n}$$

or

$$P = \frac{n}{2 + (n - 1)} + \frac{1 + (n - 1)}{2 + (n - 1)} \times c_j:$$

Substituting this into the formula for p_i, we obtain

$$\begin{array}{lll} p_{i} & = & \dfrac{n}{2+(2n-1)} + \dfrac{n}{2+(2n-1)} & \dfrac{n}{2+(n-1)} + \dfrac{1+(n-1)}{2+(n-1)} & \dfrac{x_{i}}{p-1} & c_{j} & + \dfrac{1+(n-1)}{2+(2n-1)} & c_{i} \\ \\ & = & \dfrac{n}{2+(2n-1)} & 1 + \dfrac{n}{2+(n-1)} & + \dfrac{n}{2+(2n-1)} & \dfrac{1+(n-1)}{2+(n-1)} & \dfrac{x_{i}}{p-1} & c_{j} & + \dfrac{1+(n-1)}{2+(2n-1)} & c_{i} \\ \\ & = & \dfrac{n}{2+(n-1)} + \dfrac{(1+(n-1))}{(2+(2n-1))(2+(n-1))} & \dfrac{x_{i}}{p-1} & c_{j} & + \dfrac{1+(n-1)}{2+(2n-1)} & c_{i} \end{array}$$

Thus, in equilibrium, x_i is

$$\begin{aligned} x_i &= \frac{1+n}{1+n} + \frac{1}{1+n} \overset{P}{-} \overset{P}{-} p_i \\ &= \frac{1}{1+n} + \frac{1}{1+n} & \frac{n}{2+(n-1)} + \frac{1+(n-1)}{2+(n-1)} & \overset{X_i}{-} c_j \\ & \frac{1+(n-1)}{2+(n-1)} + \frac{(1+(n-1))}{(2+(2n-1))(2+(n-1))} & \overset{X_i}{-} c_j + \frac{1+(n-1)}{2+(2n-1)} c_i \end{aligned} \overset{!}{=} \frac{f^2 + (n-1) + n - (1+n)g}{(1+n)(2+(n-1))} \\ &= \frac{f(2+(2n-1))(1+(n-1))}{(1+n)(2+(2n-1))(2+(n-1))} & \overset{X_i}{-} c_j - \frac{1+(n-1)}{2+(2n-1)} c_i \\ &= \frac{1+(n-1)}{1+n} & \overset{(1+(n-1))}{-} & \overset{(1+(n-1))}{-} & \overset{X_i}{-} & \overset{(1+(n-1))}{-} & \overset{X_i}{-} & \overset{(1+(n-1))}{-} & \overset{(1+$$

Then, ..rm i's equilibrium pro...t is

$$y_{i}(\) = \frac{1+(n-1)}{1+n} \quad \frac{1+(n-1)}{2+(n-1)} + \frac{(1+(n-1))}{(2+(2n-1))(2+(n-1))} \frac{x_{i}}{c_{j}} \quad \frac{1+n}{2+(2n-1)} c_{i} \quad \frac{x_{i}}{c_{j}} = \frac{1+(n-1)}{2+(2n-1)} c_{i} \quad \frac{x_{i}}{c_{j}} = \frac{1+(n-1)}{2+(2n-$$

From the submodularity of $f(a_m; a_w)$, $P_{i=1}^n c(m_i; w_i) = \min_{2M} P_{i=1}^n c(m_i; (m_i))$ holds. Thus, Regularity Condition 2 assures that the contents of the parenthesis is positive and $x_i > 0$ holds for all $i = 1; ...; n. \blacksquare$

Proof of Lemma 3. It is easy to see $f(m; a_w) = f(a_{m'}; a_w) = f(a_m; a_{w'}) = f(a_{m'}; a_{w'}) = 0$, since $a_m > a_{m'}$ and $a_w > a_{w'}$, and $\frac{@f}{@a} < 0$, $\frac{@f}{@a} < 0$, and $\frac{@^2f}{@a @a} = 0$. Thus, $c(m; w) + c(m^0; w^0)$

 $c(m; w^0) + c(m^0; w) \text{ holds. By letting }_w \quad c(m^0; w) \quad c(m; w) > 0 \text{ and }_{w'} \quad c(m^0; w^0) \\ c(m; w^0) > 0, \text{ we have }_w \quad w' \quad 0$

 $s_{n-2}^{\emptyset} < X_{n-1n-2} \quad (X_{n-1n-1} \quad (X_{nn-1} \quad X_{nn})). \text{ From the previous step, we know } s_{n-1}^{\emptyset} \quad s_{n-1}, \text{ and thus } \frac{\emptyset}{n-1} \quad X_{n-1n-1} \quad s_{n-1} = X_{n-1n-1} \quad (X_{nn-1} \quad X_{nn}). \text{ Thus, we have } s_{n-1}^{\emptyset} \quad s_{n-1} = X_{n-1n-1} \quad (X_{nn-1} \quad X_{nn}).$

$$S_{n-2}^{0} + r_{n-1}^{0} < X_{n-1n-2} (X_{n-1n-1} (X_{nn-1} X_{nn})) + X_{n-1n-1} (X_{nn-1} X_{nn})$$

$$= X_{n-1n-2}$$
:

This violates the stability, and contradicts with s^0 being a competitive salary. Thus $s^0_{n-2} = s_{n-2}$. Repeated applications of the same logic conclude that any competitive salary vector s^0 satis...es $s^0 = s$.

References

- [1] Bando, K. (2012), Many-to-One Matching Markets with Externalities among Firms. Journal of Mathematical Economics 48.1, 14-20.
- [2] Bando, K. (2014), A Modi..ed Deferred Acceptance Algorithm for Many-to-One Matching Markets with Externalities among Firms. Journal of Mathematical Economics 52, 173-181.
- [3] Banerjee, S., H. Konishi, and T. Sonmez (2001), Simple Coalition Formation Games, Social Choice and Welfare 18, 135-153.
- [4] Becker, G.S., (1973), A Theory of Marriage I., Journal of Political Economy 81-4, 813-846.
- [5] Bloch, F., (1996), SequexW15(o)2(F)96(o)11(r)11(m)17(a)11(t)8(i)1(r)17(.)-315(o)1411(n)11nnalitos

- [21] Herings, P.J.J., A. Mauleon, and V.J. Vannetelbosch (2020), Matching with Myopic and Farsighted Players, Journal of Economic Theory 190, 105-125.
- [22] Imamura, K., and H. Konishi, Assortative Matching with Externalities and Farsighted Players, in progress.
- [23] Kimya, M. (2021), Farsighted Objections and Maximality in One-to-one Matching Problems, University of Sydney.
- [24] Knuth, D.E., (1976), Marriages Stables, Montreal: Les Presses de l'Universite de Montreal.
- [25] Kojima, F., and U. Unver, (2008), Random paths to pairwise stability in many-to-many matching problems: a study on market equilibration, International Journal of Game Theory 36 (3), 473-488

- [33] Mumcu, A., and I. Saglam (2010), Stable One-to-One Matchings with Externalities. Mathematical Social Sciences 60.2, 154-159.
- [34] Nitzan, S. (1991), Collective Rent Dissipation, Economic Journal 101, 1522-1534.
- [35] Pycia, M., and M.B. Yenmez (2021), Matching with Externalities, University of Zurich, Department of Economics, Working Paper 392.
- [36] Ray, D., and R. Vohra (1999), A Theory of Endogenous Coalition Structures, Games and Economic Behavior 26, 286-336.
- [37] Ray, D. (2008), A Game-Theoretic Perspective on Coalition Formation, Oxford University Press, Oxford.
- [38] Ray, D., and R. Vohra (2014), "Coalition Formation," Handbook of Game Theory vol. 4, 239-326.
- [39] Rosenthal, R.W., (1972), Cooperative games in exectiveness form, Journal of Economic Theory 5, 88-101.
- [40] Sanchez-Pages, S. (2007a), Endogenous Coalition Formation in Contests, Review of Economic Design 11, 139-63.

[41]

- [46] Tullock, G., (1980), "E¢ cient Rent Seeking," Buchanan, J.M., Tollison, R.D., Tullock, G. (Eds.), Toward a Theory of the Rent-Seeking Society, Texas A&M University Press, College Station, TX, pp. 97-112.
- [47] Yi, S.-S. (1997), Stable Coalition Structures with Externalities, Games and Economic Behavior, 20(2), pp.201-237.